NEW BIRDS IN THE PHILATELIC AVIARY

Charles E. Braun, BU1364

New Birds



SOMALI OSTRICH, Struthio molybdophanes Struthionidae

Georgia, 2003 August 25, Sc#314, 40t

[Ed. note: The first appearance of this species was on a 1932 stamp of Italian Somaliland (Sc#151). That depiction is poor, so we decided to break tradition and use the much nicer looking latest issue.]

Length: 83 to 108 inches, resident. The male (shown) is black with a gray neck and white in the wings and tail. The female is brown.

Habitat: Open semi-arid plains and open woodland.

Range: Somalia to Kenya.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 1.



OLIVE IBIS, Bostrychia olivacea Threskiornithidae

Tanzania, 2015, not yet cataloged, 2000sh

Length: 28 to 30 inches, sexes alike, resident. Gray with a bushy crest, buffy neck, greenish wings and black face.

Habitat: Wetlands in dense forest.

Range: Sierra Leone to Gabon and mountains of Kenya and Tanzania.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. *Handbook of the Birds*

of the World, Volume 1.



AFRICAN SWAMPHEN, Porphyrio madagascariensis Rallidae

Burundi, 1979 May 30, Sc#553, 10fr

Length: 15 to 20 inches, sexes similar, resident. Bronzy-green or blue-green back and scapulars and purplish below with a turquoise neck and breast and red bill, shield, and legs.

Habitat: Reed beds, marshes, and flooded grasslands.

Range: Egypt, sub-Saharan Africa, and Madagascar.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 3.



BURCHELL'S COURSER, Cursorius rufus Glareolidae

Namibia, 2015, not yet cataloged, postcard rate

Length: 8 to 9 inches, sexes alike, resident. Pale rufous-brown above and light rufous-brown below with a blue-gray hindcrown, white superciliary, and black stripes behind the eye.

Habitat: Open plains with short grass.

Range: Southwestern Angola to South Africa.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. *Handbook of the Birds of the World, Volume 3*.



DOUBLE-BANDED COURSER, Smutsornis africamus Glareolidae

Namibia, 2015, not yet cataloged, Zone B

[Ed. note: We use Clements as our authority for taxonomy and naming. However, other

authorities note Smutsornis as a junior synonym of Rhinoptilus.]


Length: 8 to 10 inches, sexes alike, resident. Scaly Buff-brown above and crown and cinnamon below with two black breast bands.

Habitat: Stony and gravelly semi-desert.

Range: Ethiopia to South Africa

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 3.



THREE-BANDED COURSER, Rhinoptilus cinctus Glareolidae

Namibia, 2015, not yet cataloged, Zone C

Length: 10 to 12 inches, sexes alike, resident. Scaly dark brown above and breast and white below with chestnut and black bands.

Habitat: Open sandy clearings, scrubby grassland, and woodlands with bare ground.

Range: Sudan to Zambia.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 3.



BRONZE-WINGED COURSER, Rhinoptilus chalcopterus Glareolidae

Namibia, 2015, not yet cataloged, Zone A

Length: 10 to 12 inches, sexes alike, inter-African migrant. Grayish-bronze above and

white below with a pale brown neck and a dark band on the lower breast.

Habitat: Dry savanna, woodland, and thick bush.

Range: Senegal to Sudan and South Africa.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 3.



MOURNING COLLARED DOVE, Streptopelia decipiens Columbidae

Mozambique, 2015 June 15, not yet cataloged, 46m


Length: 11 inches, sexes alike, resident. Brown above and rosy gray below with a gray head and a white-margined black hindcollar.

Habitat: Thorny acacia, woods, thickets, and arid savanna.

Range: Senegal to Somalia, Zambia, and Angola.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 4.



COMMON BRONZEWING, Phaps chalcoptera Columbidae

Mozambique, 2015 June 15, not yet cataloged, 46m

Length: 11 to 14 inches, sexes similar, resident. Scaly olive-brown above and brownish pink below with a yellow forehead and iridescent wing spots.

Habitat: Woodlands.

Range: Australia and Tasmania.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 4.



SJOSTEDT'S OWLET, Glaucidium sjostedti Strigidae

Central African Republic, 2015 October 21, not yet cataloged, 900fr

Length: 8 to 10 inches, sexes alike, resident. Deep chestnut above and barred cinnamon

below with barred head, wings, and tail and white chin and brows.

Habitat: Primary lowland forest.

Range: Cameroun to Gabon, Southern Central African Republic, and central Zaire.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 5.



BUFF-FRONTED OWL, Aegolius harrisii Strigidae

Uruguay. 2015 September 30, not yet cataloged, \$15

Length: 8 inches, sexes alike, resident. White-spotted chocolate-brown above and vellowish-buff below with a buff forehead.

Habitat: Open humid forest.

Range: Northwestern Venezuela to north-central Peru, and eastern Bolivia to

eastern Brazil and Uruguay.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5.



DIMORPHIC DWARF-KINGFISHER, Ceyx margarethae Alcedinidae

Philippines, 2007 November 15, Sc#3128b, 20p

Length: 6 inches, sexes alike, resident. Pale blue (pale morph) or dark blue (dark morph) above and rich orange below with a white chin and ear patch and red bill.

Habitat: Rainforest, monsoon forest, and thick vegetation along streams.

Range: Central and southern Philippines.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 6.



NORTH SOLOMONS DWARF-KINGFISHER, Ceyx meeki Alcedinidae

Solomon Islands, 2001 February 1, Sc#910, \$4

Length: 6 inches, sexes alike, resident. Dark blue above, paler in the center, and vellowish below with a white chin and black bill.

Habitat: Rainforest and thick vegetation along streams.

Range: Western and central Solomon Islands.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. *Handbook of the Birds of the World, Volume 5*.



PACIFIC KINGFISHER, Todiramphus sacer Alcedinidae

Niuafo'ou, 1983 November 15, Sc#38, 32c

Length: 10 inches, sexes alike, resident. Blue-green above and white or rufous below with a rufous superciliary and black eye-stripe.

Habitat: Mainly coastal.

Range: Polynesia.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World,

Volume 6.



MELANESIAN KINGFISHER, Todiramphus tristrami Alcedinidae

Solomon Islands, 2004 November 15, Sc#985b. \$7.50

Length: 9 to 10 inches, sexes alike, resident. Blue and green above and rufous below with a rufous front and a white throat.

Habitat: Coasts and lowland forests.

Range: New Britain to the Solomon Islands and Vanuatu.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 6.



SPOT-BILLED TOUCANET, Selenidera maculirostris Ramphastidae

Argentina, 2015 September 22, not yet cataloged, \$7

Length: 13 to 14 inches, resident. The male (shown) is green above (and eye patch) and black below (and neck and crown) with gold-yellow ear tufts, rusty thighs, and red undertail. The female is red with green back, wings, and eye patch and a blue undertail.

Habitat: Subtropical forest.

Range: Southeastern Brazil to eastern Paraguay.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 7.



ROCK KESTREL, Falco rupicolus Falconidae

Namibia, 1999 May 18, Sc#937, \$1.60

Length: 12 to 14 inches, sexes similar, resident. Brown, paler and

streaked below with a gray head and black-barred tail.

Habitat: Grassland, scrub, and open woodland.

Range: Northern Angola to southern Tanzania and South Africa.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 2.



NORTH ISLAND SADDLEBACK, Philesturnus rufusater Callaeidae

New Zealand, 1986 April 23, Sc#770A, \$4

Length: 10 inches, sexes alike, resident. Glossy black with a dark orange or red-brown saddle.

Habitat: Forest.

Range: Small islands off the North Island, New Zealand.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 14.



SOUTH ISLAND SADDLEBACK, Philesturnus carunculatus Callaeidae

New Zealand, 1962 October 3, Sc#B64, 3p

Length: 10 inches, sexes alike, resident. Glossy black with a dark orange or red-brown

saddle.

Habitat: Forest.

Range: Islands off northeastern and southwestern South Island, New Zealand.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World,

Volume 14.



SOOTY MYZOMELA, Myzomela tristrami Meliphagidae

Solomon Islands, 2015 September 3, not yet cataloged, \$12

Length: 5 inches, sexes similar, resident. All black.

Habitat: Forests, plantations, and gardens.

Range: Eastern Solomon Islands.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds

of the World, Volume 13.



YELLOW-VENTED MYZOMELA, Myzomela eichhorni Meliphagidae


Solomon Islands, 2015 September 3, not yet cataloged, \$12

Length: 5 inches, sexes similar, resident. Dark olive-gray above and yellowolive below with a scarlet rump and throat.

Habitat: Forest, forest edges, plantations, and gardens.

Range: West-central Solomon Islands.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds





$\textbf{NOISY MINER}, \textit{Manorina melanocephala} \ \textbf{Meliphagidae}$

Solomon Islands, 2015 September 3, not yet cataloged, \$12

Length: 10 to 11 inches, sexes alike, resident. Gray with a white belly and patterned head.

Habitat: Dry open forests and woodlands.

Range; Eastern Australia and Tasmania.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds

of the World, Volume 13.



ASIAN PARADISE FLYCATCHER, Terpsiphone paradisi Monarchidae

Russia, 1981 August 20, Sc#4973, 10k

Length: 7 inches plus a 4- to 7-inch tail streamer, sexes similar but the female lacks the tail streamer, migratory. Deep rufous above and white or pale gray below with a black head and crest.

Habitat: Forest, shrubs, and plantations.

Range: Afghanistan to Sri Lanka, peninsular Malaysia and Myanmar. Winters from Pakistan to Sri Lanka and the Malay Peninsula.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. *Handbook of the Birds of the World, Volume 11*.



BLYTH'S PARADISE-FLYCATCHER, *Terpsiphone affinis* Monarchidae Malaysia, 1965 September 9, Sc#26, \$5

Length: 7 inches plus a 4- to 7-inch tail streamer, sexes similar but the female lacks the tail streamer, resident. Deep rufous above and gray below with a black head and crest. The bird on the right is a white morph.

Habitat: Forests, shrubs, and plantations.

Range: Southern China to the Lesser Sundas.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 11.



the World, Volume 11.

ISLAND MONARCH, Monarcha cinerascens Monarchidae

Solomon Islands, 2015 September 3, not yet cataloged, \$12

Length: 6.5 to 7.5 inches, sexes alike, resident. Dull gray with a rufous belly.

Habitat; Forests and coastal scrub.

Range: Lesser Sundas and Sulawesi to the Solomon Islands.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of



BLACK-WINGED MONARCH. Monarcha frater Monarchidae

Solomon Islands, 2015 September 3, not yet cataloged, \$12

Length: 6 to 8 inches, sexes similar, resident. Pale gray with a black face, wings, back and tail and orange-rufous belly.

Habitat: Forests and mangroves.

Range: New Guinea, northern Australia, and the northern Solomon Islands.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 11.



WHITE-CAPPED MONARCH, Monarcha richardsii Monarchidae

Solomon Islands, 2015 September 3, SS not yet cataloged, \$40

Length: 7 inches, sexes alike, resident. Glossy bluish-black with a chestnut belly and white nape.

Habitat: Semi-evergreen forests.

Range: New Georgia Islands and the western Solomon Islands.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 11.