NEW BIRDS IN THE PHILATELIC AVIARY Charles E. Braun, BU1364

Corrections

During the assembly of this column for Vol. 64(4), the editor mistakenly mixed up the images and descriptions for the Burchell's, Three-banded, and Bronze-winged Coursers. Because the name of the bird on the stamp itself is easily discerned, you can match the correct description to each one.





BRAZILIAN TEAL, Amazonetta brasiliensis Anatidae

St. Martin, 2016 May 25, not yet cataloged, 25c

Length: 14 to 16 inches. The male (shown) is dark brown above and speckled rufous below with a red bill and gray face and throat. The female has a dark bill and white face and throat.

Habitat: Inland ponds and small lakes in densely wooded areas.

Range: Eastern Colombia to northern Argentina. Southern birds winter closer to the equator.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 1.



EASTERN CHANTING GOSHAWK, Melierax poliopterus Accipitridae

Djibouti, 2016 May 5, not yet cataloged, 280fr

Length: 18 to 20 inches, sexes alike, resident. Ash gray with a barred white belly and a black tail.

Habitat: Dry thorn woodland and savanna.

Range: Somalia to northern Tanzania.

Reference: del Hovo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 2.



SHARP-TAILED GROUSE, Tympanuchus phasianellus

Phasianidae

Previously on a Saskatchewan Conservation stamp. Canada, 2016 July 12, not yet cataloged, (85c)

Length: 16 to 19 inches, resident. The male is speckled brown with a short pointed tail and an inflatable purple neck sac.

The female lacks the neck sac.

Habitat: Open areas and forests.

Range: Alaska to Hudson Bay and Wyoming.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 2.



SLATY-BREASTED WOOD-RAIL Aramides saracura Rallidae

St. Martin, 2016 May 25, not yet cataloged, 125c

Length: 14 to 15 inches, sexes alike, resident. Brown above and gray head and below with a rufous nape and mostly green bill.

Habitat: Swampy areas in forests and woodland.

Range: Southeastern Brazil to eastern Paraguay.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds

of the World, Volume 3.



SHORE PLOVER, Thinornis novaeseelandiae Charadriidae

Togo, 2016 May 20, not yet cataloged, 3000fr SS

Length: 8 inches, sexes similar, resident. Gray-brown above and white below with a black face, neck, and throat and a white-edged gray-brown crown.

Habitat: Rocky shores.

Range: Rangatira Island, Chatham Islands, New Zealand.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 3.\\



BAIRD'S SANDPIPER, Calidris bairdii Scolopacidae

St. Martin, 2016 May 25, not yet cataloged. 200c

Length: 6 to 7 inches, sexes alike, migratory. Brown above and white below with a buffy breast.

Habitat: Low mountain tops, river terraces, and coastal barren areas.

Range: Breeds from northern Alaska to western Greenland. Winters in western and southern South America.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds

of the World, Volume 3.



SCALED DOVE, Columbina squammata Columbidae

St. Martin, 2016 January 6, not yet cataloged, 500c

Length: 7 to 9 inches, sexes alike, resident. Brown above and grayish pink below with a white throat and a patch of spots on the wings.

Habitat: Arid tropical savanna and thickets.

Range: Colombia to French Guiana; Central and eastern Brazil and adjacent Argentina and Paraguay.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 4.



SULAWESI GROUND-DOVE, Gallicolumba tristigmata Columbidae

Sierra Leone, 2016 January 28, not yet cataloged, 6000Le

Length: 13 to 14 inches, sexes alike, resident. Olive-brown above and gray below with a bright golden forehead, gold lower breast, and whitish belly.

Habitat: Lowland primary forest.

Range: Sulawesi, Indonesia.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 4.



WHITE-CRESTED TURACO, Tauraco leucolophus Musophagidae

Liberia, 2016 February 20, not yet cataloged, \$100

Length: 16 inches, sexes alike, resident. Violet-blue above and grass green below with a white crest and red eye-ring on a black face.

Habitat: Riverine gallery forest and wooded savannas.

Range: Southeastern Nigeria to western Kenya.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 4.



LARGE HAWK-CUCKOO, Hierococcyx sparverioides Cuculidae

Sierra Leone, 2016 January 28, not yet cataloged, MS/4 (L margin)

Length: 15 to 16 inches, sexes alike, migratory. Brown above and barred white below with a rufous or rufous-streaked breast.

Habitat: Deciduous evergreen woods.

Range: Breeds from the Himalayas and China to Borneo and the Philippines. Winters from

eastern India to Borneo and the Philippines.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World,

Volume 4.



ASIAN EMERALD CUCKOO, Chrysococcyx maculatus Cuculidae

Sierra Leone, 2016 January 28, not yet cataloged. 6000Le (Cap. C. cupreus)

Length: 7 inches, migratory. The male is green with a green-barred white belly; the female (shown) is green with a light rufous crown and nape and barred pale throat. Habitat: Dense forest.

Range: Breeds from central China to the Andaman Islands and central Vietnam. Winters from southern China to Malaysia.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 4.

This set also includes a stamp captioned *Cacomantis merulinus* (Plaintive Cuckoo) that would be a new species, but the illustration shows a different bird, unidentified but definitely not a cuckoo.



LEWIS'S WOODPECKER, Melanerpes lewis Picidae


Central African Republic, 2016 June 20, not yet cataloged, 750fr (on left)

Length: 10.5 to 11.5 inches, sexes alike, migratory. Blackish above and pinkish below with a crimson face, black throat, and silver-gray neck and breast.

Habitat: Breeds in open woodlands. Winters in oak woodland and orchards.

Range: Breeds from southern British Columbia to North Dakota, Arizona, and New Mexico. Winters in the southwestern U.S.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 7.



GILDED FLICKER, Colaptes chrysoides Picidae

Mozambique, 2016 January 15, not yet cataloged, 66m

Length: 10 to 12 inches, resident. The male (shown) is black-barred brownish above and black-spotted whitish below with a rufous crown and mustache and white rump. The female lacks the mustache.

Habitat: Arid areas with large cacti and yucca plants.

Range: Southeastern California to northern Mexico and Baja California.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 7.



PLAIN PARAKEET. Brotogeris tirica Psittacidae

St. Martin, 2016 January 6, not yet cataloged, 500c

Length: 9 inches, sexes alike, resident. Green with bronzy green shoulders.

Habitat: Rainforests, open areas, parks, and plantations.

Range: Coastal southeastern Brazil.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of

the World, Volume 4.



RED-RUFFED FRUITCROW, Pyroderus scutatus Cotingidae

Colombia, 2016 July 29, label on SS

Length: 17 to 18 inches, sexes similar, but female is smaller, resident. Black above and reddish below.

Habitat: Forest and forest borders.

Range: Colombia, central Peru, eastern Venezuela, and southeastern Brazil. Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 9.



CAVE SWALLOW, Petrochelidon fulva Hirundinidae

Kosovo, 2010 September 23, Sc#169, €1

Length: 5 inches, sexes alike, migratory. Glossy blackish above and cinnamon and white below with white streaks on the mantle and a cinnamon rump.

Habitat: Open areas often near water.

Range: Breeds from southern New Mexico and Texas to southern Mexico and the West Indies. U.S. birds winter in Mexico and Central America.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 4.



RED-BREASTED NUTHATCH, Sitta canadensis Sittidae

Previously on a local stamp from Eynhallow, Scotland. United States, 2016 August 4, not yet cataloged, (47c)

Length: 5 inches, sexes similar, resident. Gray above and rufous below with a white face and a black crown and eye-stripe.

Habitat: Temperate forests.

Range: Southern Alaska to Newfoundland, western North Carolina, and New Mexico.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 13.



BROWN CREEPER, Certhia americana Certhiidae

Central African Republic, 2016 June 20, not yet cataloged, 750fr

Length: 5 inches, sexes alike, migratory. White-streaked brown above and whitish below.

Habitat: Coniferous forests.

Range: Breeds from southern Alaska to Newfoundland and Nicaragua. Winters in the

breeding area except northern Canada.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 13.



BALLMAN'S MALIMBE, Malimbus ballmanni Ploceidae

Sierra Leone, 2016 May 27, not yet cataloged, 6000Le

Length: 6 to 7 inches, resident. The male is black with a yellow nape, breast, and

undertail. The female (shown) lacks the yellow nape.

Habitat: Lowland evergreen forest.

Range: Extreme eastern Sierra Leone and adjacent Liberia.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 15.



BLACK-HEADED WEAVER, Ploceus melanocephalus Ploceidae

Tanzania, 2016 January 28, not yet cataloged, 6000sh SS (Cap: P. intermedius)

Length: 6 inches, resident. The male (shown) is greenish-yellow above and yellow or chestnut below with a black head and yellow nape. The female is greenish above and white below with a pale yellow breast.

Habitat: Tall grass in damp areas near water.

Range: Mauritania to Kenya and Zambia.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 15.



CHESTNUT-EARED BUNTING, Emberiza furcata Emberizidae

Japan, 2016 August 10, not yet cataloged, 82¥

Length: 6 inches, migratory. The male (shown) is brown above and pale below with a gray head, chestnut ears, black mustache and neck stripe, and rufous breast band. The female is paler with no black.

Habitat: Scrubby hillsides and open areas.

Range: Breeds from eastern Mongolia to Korea and Japan. Winters from Japan to the Himalayas and Myanmar.

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 16.



PATAGONIAN SIERRA-FINCH, *Phrygilus patagonicus* Emberizidae St. Vincent (Bequia), 2016 March 1, not yet cataloged, \$3.15

Length: 6 inches, resident. The male (shown) is dusky above and yellow below with a blue-gray head and russet nape. The female lacks the russet nape and has streaks below. Habitat: Forest, roadsides, and clearings.

Range: Chile and western Argentina.

Reference: del Hoyo, J., A. Elliott. and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 16.