NEW BIRDS IN THE PHILATELIC AVIARY

Charles E. Braun, BU1364

Corrections

In Biophilately Vol. 60(3), BUFF-STREAKED LORY should have been BLUE-STREAKED LORY, Eos reticulata.

New Birds



PLUMED WHISTLING DUCK, Dendrocygna eytoni Anatidae

Previously on an Australian Wildlife Conservation stamp Australia, 2012, not yet cataloged, \$2.35

Length: 16 to 18 inches, sexes alike, disperses. Buffy head, neck and breast,

black-streaked chestnut abdomen and long flank plumes.

Habitat: Grassland, meadows and plains

Range: Breeds in coastal eastern Australia; disperses to northern and east-central Australia

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 1



BEAUDOUIN'S SNAKE-EAGLE, Circaetus beaudouini Accipitridae

Uganda, 2012, not yet cataloged, 3400s

Length: 24 inches, sexes alike, resident. Brown above and barred white below

with a brown breast, pale chin and three dark bars in the tail.

Habitat: Open woodland and tree savanna

Range: Mauritania to northwestern Kenya

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of

the World, Volume 2



DAMARA TERN, Sternula balaenarum Laridae

Previously on postal stationery from Southwest Africa

Namibia, 2012, not yet cataloged, \$20

Length: 9 inches, sexes alike, disperses. Light gray above and white below with a black crown and bill.

Habitat: Seacoast

Range: Breeds in Namibia and South Africa; disperses to Nigeria

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 3



AUSTRALIAN MASKED-OWL, Tyto novaehollandiae Tytronidae

United Nations, 2012, not yet cataloged, €0.70

Length: Male is 13 to 17 inches; female is 17 to 23 inches, resident. Blackish-brown above with a rufous wash and white speckles and pale rufous below with dark speckles with a dark chestnut facial disc. A pale morph occurs that is white below with a white facial disc.

Habitat: Tall open forest

Range: Australia except dry central regions and Tasmania

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5



RUFOUS TWISTWING, Cnipodectes superrufus Tyrannidae

Guinea, 2011, not yet cataloged, 40,000fr

Length: 6 to 7 inches, sexes alike, resident. Bright rufous overall.

Habitat: Lower growth of humid, deciduous forest and secondary woodland

Range: Southwestern Amazonia in Brazil, Bolivia, and Peru

Reference: Lane, D. G., G. P. Servat, T. Valqui, and F. R. Lambert. The Auk 124(3): 762-

772, 2007



BALD PARROT, Pyrilia aurantiocephala Psittacidae

Guinea, 2011, not yet cataloged, 40,000fr (upper left)

Length: 9 inches, sexes alike, resident. Green with a bald brownish-orange head and red

underwings.

Habitat: Lowland forest

Range: East-central Amazonian Brazil

Reference: Gaban-Lima, R., M. A. Raposo, and E. Hoefling. The Auk, 119 (3): 815-818, 2002



CAMIGUIN HANGING-PARROT, Loriculus camiguinensis Psittacidae

Guinea, 2011, not yet cataloged, 4×10,000fr (left margin)

Length: 5 inches, sexes similar, resident. Green above and yellow-green below with a blue face, throat, and undertail and red crown and upper tail

Habitat: Lowland trees and bushy areas

Range, Camiguin Island, Philippines

Reference: Telio, J.G., J.F. Degner, J. M. Bates, and D. E. Willard. Fieldiana, Zoology 106,

49-57, 2006



GORGETED PUFFLEG, Eriocnemis isabellae Trochilidae

Guinea, 2011, not yet cataloged, 4×10,000fr (center left margin)

Length: 3.5 to 4 inches, resident. The male (shown on the stamp) is dark with an iridescent green and violet-blue throat patch and a white belly and thighs; the female lacks the throat patch.

Habitat: Cloud forest

Range: Southwestern Colombia

Reference: Cortes-Diago, A., L.A. Ortega, L. Mazariegos, and A. A. Weller. Ornitologia Neotropical 18: 161-

170, 2007



VIOLET WOODHOOPOE, Phoeniculus damarensis Phoeniculidae

Namibia, 2012, not yet cataloged, registered

Length: 14 to 15 inches, sexes similar, resident. Dark with a coppery purple iridescence with white spots on the wings and outer tail and red bill and legs.

Habitat: Dry acacia shrub and thorn bush

Range: Southern Ethiopia , Kenya, western Angola and northwestern Namibia Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World. Volume 6



DAMARA RED-BILLED HORNBILL, Tockus damarensis Bucerotidae

Namibia, 2012, not yet cataloged, Non-Standard rate

Length: 14 inches, sexes similar but the female has less black on the lower mandible, resident. Black above and white face, below and wing spots with a red and black bill and fleshy skin around the eyes.

Habitat: Open savanna and woodland Range: Northwestern and central Namibia

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 6



WHITE-BELLIED CINCLODES, Cinclodes palliatus Furnariidae

Peru, 2011, not yet cataloged, 10s

Length: 9 inches, sexes alike, resident. Brown above and gleaming white below with a grayish crown, black lore and a white wing band.

Habitat: Bogs and short grass Range: High Andes of central Peru

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 8



CUBAN PEWEE, Contopus caribaeus Tyrannidae

Cuba, 2011, not yet cataloged, 5c

Length:6 to 7 inches, sexes similar, resident. Olive-brown above and beige-gray below with an olive-gray crown and buffy mustard-yellow belly.

Habitat: Forest, forest edges, swamps, mangroves, brushy scrub and tree plantations

Range: Cuba and the Bahamas

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 9



CHESTNUT-BACKED CHICKADEE, Poecile rufescens Paridae

Guinea-Bissau, 2011, not yet cataloged, 3500fr


Length: 5 inches, sexes alike, resident. Black cap and chin, white face and belly and chestnut back and sides.

Habitat: Dense conifer forests

Range: Southern Alaska to coastal California

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 12



CARP'S TIT, Parus carpi Paridae

Namibia, 2012, not yet cataloged, 5c


Length: 6 inches, sexes similar but the female is duller, resident. Black with white in the wings and outer tail.

Habitat: Savanna, acacia woodland and thickets

Range: Southwestern Angola and northern and western Namibia

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 12



BARE-CHEEKED BABBLER, Turdoides gymnogenys Timaliidae

Namibia, 2012, not yet cataloged, \$100

Length: 10 inches sexes alike, resident. Brown above and pale below with a pale gray crown and barred black cheeks (should be called bar-cheeked rather than bare-cheeked).

Habitat: Undergrowth of woodland and thickets

Range: Southwestern Angola and northwestern Namibia

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 12



BLACK-COLLARED STARLING, Gracupica nigricollis Sturnidae

Surinam, 2008, 1376, label

Length:11 inches, sexes alike, resident. Black above and white below with a white head, broad black collar and bare yellow skin around the eye.

Habitat: Open areas and dry stunted forest

Range: Burma to southeastern China, Vietnam, and peninsular Thailand

Reference: del Hoyo, J., A. Elliott, and J. Sargatal, Eds. Handbook of the Birds of the

World, Volume 14