### NEW BIRDS IN THE PHILATELIC AVIARY Charles E. Braun -BU 1364

#### **New Birds**

[Continued from Vol. 55 (1)]

#### **SNOWY-CROWNED ROBIN-CHAT,** Cossypha niveicapilla

Muscicapidae (142.161)

Staffa, 1982, uncataloged, 20 pence

Length: 9 inches, sexes alike, resident. Black face and above and orange

below and collar with a white crown.

Habitat: gallery forest and dense thickets.

Range: Senegal to Ethiopia and western Tanzania.

Reference: Sinclair, I and P.Ryan. Birds of Africa South of the Sahara.





**DUSKY SUNBIRD,** Cinnyris fusca Nectariniidae (166.089)

Namibia, 2005, not yet cataloged, \$3.40

Length: 4 inches, resident. The male (shown on the stamp) is black with a white belly and orange pectoral tufts; the female is light gray-brown above and off-white below.

Habitat: dry country.

Range: Southwestern Angola to western South Africa. Reference: Cheke, R. A. and C. F. Mann. Sunbirds.



**RED-WINGED PYTILIA,** *Pytilia phoenicoptera* Estrildidae (193.011) Staffa, 1982, uncataloged, 60 pence

Length: 5 inches, sexes similar but the female is paler, resident. Gray with red wings and tail.

Habitat: semiarid savanna, open woodland and thickets.

Range: Senegal to western Ethiopia.

Reference: Sinclair, I and P.Ryan. Birds of Africa South of the Sahara.

RAROTONGA STARLING, Aplonis cinerascens Sturnidae

(190.021) Cook Islands, 2005, not yet cataloged, \$1.15

Length: 8 inches, sexes alike, resident. Gray with white undertail.

Habitat: undisturbed montane forest. Range: Raratonga, Cook Islands.

Reference: Pratt, H. D., P.L. Bruner and D. G. Berrett. The Birds of Hawaii and the Tropical Pacific.



YELLOW-RUMPED WARBLER, *Dendroica coronata* Parulidae (199.019) British Virgin Islands, 2005, not yet cataloged, 15 cents The 60-cent stamp in this set shows the Cape May Warbler which was written up in *Biophilately* 54 #2.

Length: 6 inches, sexes similar but the female is duller, migratory. Gray above and streaked white below with yellow patches on the crown. The stamp shows the Myrtle Warbler, *Dendroica coronata coronata*. The Audubon Warbler has a yellow throat.

Habitat: breeds in woodlands; winters in many habitats.

Range: Breeds from Alaska to northern Mexico and Maryland; winters from the southern U.S. to

Panama and the West Indies.

Reference: Curson, J., D. Quinn and D. Beadle. Warblers of the Americas.

**BLACK-AND-WHITE TANAGER,** *Conothraupis speculigera* Thraupidae (201.017)

Eynhallow, 1982, uncataloged, 10 pence

Length: 6.5 inches, resident. The male (shown on the stamp) is black with a white belly nad wing spot; the female is olive above and pale yellow below with a streaked breast..

Habitat: deciduous woodland, arid scrub and borders of humid forest.

Range: Western Ecuador and northern Peru.

Reference: Ridgely, R. S. and G. Tudor. The Birds of South America,

Volume I.





**CRESTED OWL,** *Lophstrix cristata.* Strigidae (078.113) Surinam, 2005, not yet cataloged, \$1.80

Length: 15 to 17 inches, sexes alike, resident. Brown, paler below with white wing spots and brows that extend to ear tufts.

Habitat: forest. Range: Southern Mexico to northern Bolivia.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of

the World, Volume 5.



#### **BLACK-THROATED GREEN**

**WARBLER**, *Dendroica virens* Parulidae (199.022)

Staffa, 1982, uncataloged, 1 pound

Length: 5 inches, sexes similar but the female is paler and duller, migratory. Olive green above and white below with a black throat and flank streaks and a yellow facial pattern.

Habitat: open forests.

Range: Breeds in central and eastern Canada, south through the Appalachians;

winters in southern Florida, the West Indies and Central America.

Reference: Curson, J., D. Quinn and D. Beadle. Warblers of the Americas.

**YELLOWBILL**, *Ceuthmochares aereus* Cuculidae (076.053) Staffa, 1982, not cataloged, 75 pence

Length: 13 inches, sexes alike, resident. Glossy above and gray below with a vellow bill.

Habitat: forest.

Range: Gambia to Angola and southern Sudan; Eritrea to eastern South

Africa.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the

Birds of the World, Volume 4.





**STRIPED OWL,** *Pseudoscops clamator* Strigidae (078.182) Surinam, 2005, not yet cataloged, \$0.05

Length: 12 to 15 inches, sexes alike, resident. Buffy, paler below with heavy black stripes and a black-bordered facial disk.

Habitat: forest, savannah, marshes and scrub.

Range: Southern Mexico to northern Argentina.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5.



WHIP-POOR-WILL, Caprimulgus vociferus Caprimulgidae (083.033) Staffa, 1982, not cataloged, 10 pence

Length: 9 to 11 inches, migratory. The male (shown on the stamp) is streaked brown above and white-spotted brownish below with white outer tail tips; the female lacks the white-tipped tail.

Habitat: forest and open woodland.

Range: Breeds in the eastern U. S. and bordering Canada and from Arizona to Honduras; winters in southeastern U.S. and Mexico to Honduras.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5.





**SCALE-THROATED HERMIT,** *Phaethornis eurynome* Trochilidae (086.021) [above left] Eynhallow, 1981, not cataloged, 10 pence

Length: 6 inches, sexes similar, resident. Olive above and grayish below with a scaly throat.

Habitat: humid forests.

Range: Southeastern Brazil to eastern Paraguay.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5.

**RUFOUS-CRESTED COQUETTE,** *Lophornis delattrie* (probable) Trochilidae (086.078) [right] Bernera, 1981, not cataloged, 10 pence

Length: 3 inches, resident. The male (shown on the stamp) is bronzy-green with a rufous crest, green cheek tufts and a white rump band; the female lacks the crest and tufts.

Habitat: forest edges.

Range: Southwestern Costa Rica tonorthern Bolivia.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5.




**FESTIVE COQUETTE,** *Lophornis chalybeus* Trochilidae (086.080) Bernera, 1981, not cataloged, 75 pence

Length: 3 inches, resident. The male (shown on the stamp) is green with a white rump band, stiff head feathers and cheek tufts; the female lacks the tufts.

Habitat: humid forest, second growth and cerrado.

Range: Southern Venezuela to central Bolivia; southeastern Brazil.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5.



**BLACK-BELLIED THORNTAIL,** *Discosura langsdorffi* Trochilidae (086.085) Staffa, 1981, not cataloged, 10 pence

Length: The male is 5 inches and the female 3 inches, resident. The male (shown on the stamp) is coppery green with a white rump band and black belly, wings and long tail; the female lacks the black belly and long tail.

Habitat: humid lowland forest and edges.

Range: Western Amazonia and coastal east-central Brazil.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds.

Handbook of the Birds of the World, Volume 5.



WHITE-VENTED PLUMELETEER, *Chalybura buffonii*Trochilidae (086.179) Bernera, 1982, not cataloged, 75 pence

Length: 4 to 5 inches, resident. The male (shown on the stamp) is green with varying amount of blue below, brown wings and white belly; the female is pale gray below.

Habitat: forest and forest edges.

Range: Eastern Panama to northwestern Venezuela.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5.



**BRONZY INCA,** *Coeligena coeligena* Trochilidae (086.221) Staffa, 1981, not cataloged, 40 pence

Length: 4.5 inches, sexes similar, resident. Bronzy-brown above and rufous-brown below with a spotted throat and neck. Habitat: humid montane cloudforest.

Range: Venezuela to southeastern Bolivia.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds.

Handbook of the Birds of the World, Volume 5.



**BLACK INCA,** *Coeligena prunellei* Trochilidae (086.223) Bernera, 1982, not cataloged, 60 pence

Length: 5 inches, sexes similar, resident. Purplish-black with a long, straight black bill.

Habitat: humid montane forest.

Range. Andes of Colombia.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds.

Handbook of the Birds of the World, Volume 5.



#### **BEARDED MOUNTAINEER,** Oreonympha nobilis

Trochilidae (086.269) Bernera, 1982, not cataloged, 40 pence

Length: 6 inches, resident. The male (shown on the stamp) is brown above and white below with a blue and purplish beard; the female is greenish above with a green throat and patchier beard.

Habitat: montane woodland and scrub.

Range: Central and southern Peru.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds.

Handbook of the Birds of the World, Volume 5.



#### **GLOW-THROATED HUMMINGBIRD,** Selasphorus ardens

Trochilidae (086.332)

Staffa, 1982, not cataloged, 1 pound

Length: 3 inches, resident. The male (shown on the stamp) is bronzy-green above with a pinkish-red gorget, white collar and buffy underparts; the female has a speckled buff throat.

Habitat: clearings and forest borders.

Range: West-central Panama.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook

of the Birds of the World, Volume 5.



**BLACK-EARED FAIRY,** *Heliothryx aurita* Trochilidae (086.294) Staffa, 1982, uncataloged, 25 pence

Length: 4 to 5 inches, sexes similar, resident. Green above and white below with black and purple face patches and dark wings and tail.

Habitat: wet lowland forest and forest edges.

Range: Southeastern Colombia to southeastern Brazil and central Bolivia. Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 5.



# **RUFOUS-HEADED GROUND-ROLLER,** *Atelornis crossleyi* Brachypteraciidae (094.004) Staffa, 1982, uncataloged, 2 pounds

Length 10 to 11 inches, sexes alike, resident. Dark green above and deep rufous head and below with a white-streaked black throat.

Habitat: humid rainforest. Range: Eastern Madagascar.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 6.




**RED-CROWNED BARBET,** *Megalaima rafflesi* Capitonidae (101.010) Singapore, 2005, not yet cataloged, \$1


Length: 10 inches, sexes alike, resident. Green, paler below, with a red crown and spot below the eye and black and yellow face.

Habitat: forest.

Range: Southern Myanmar to Borneo.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 7.





**GOLD-WHISKERED BARBET,** *Megalaima chrysopogon* Capitonidae (101.009) [on the left] Eynhallow, 1982, not cataloged, 2 pounds

Length: 12 inches, sexes similar, resident. Green, paler below with a red forehead, red hindcrown and yellow forecrown and malar area.

Habitat: forest and forest clearings.

Range: Myanmar to Borneo.


Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 7.


**FINE-SPOTTED WOODPECKER,** *Campethera punctuligera* Picidae (104.060) [on the right] Staffa,1982, not cataloged, 2 pounds

Length: 9 inches, resident. The male (shown on the stamp) is green above and white or yellow below with fine spots and a red crown and malar stripe; the female lacks the red malar stripe and forecrown. Habitat: wooded savanna.

Range: Senegal to Sudan.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 7.





**BANDED BROADBILL**, *Eurylaimus javanica* Eurylaimidae (105.007) [on the left] Eynhallow, 1982, not cataloged, 2 pounds

Length: 9 inches, resident. The male (shown on the stamp) has a brown back with yellow patches, purplish pink underparts

and purple-red head; the female is similar with a grayish wash on the breast.

Habitat: forest.

Range: Southeastern Myanmar to Vietnam and Java.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 8.

**BLACK-BACKED ANTSHRIKE,** *Sakesphorus melanonotus* Thamnophilidae (109.013)[on the right] Grunay, 1982, not cataloged, 2 pounds

Length: 6 inches, resident. The male (shown on the stamp) is black with a white belly, wing edgings and tail tips; the female is brown above and pale buff below with a dark brown crown and a rufous tail.

Habitat: deciduous forest.

Range: Caribbean slope in Colombia and northwestern Venezuela.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the Birds of the World, Volume 8.



#### BARRED ANTTHRUSH, Chamaeza mollissima

Formicariidae (110.011)

Bernera, 1982, not cataloged, 60 pence

Length: 8 inches, sexes alike, resident. Chestnut brown above and finely barred below.

Habitat: cloudforest and humid montane forest.

Range: Colombia to central Bolivia.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Hand-

book of the Birds of the World, Volume 8.

**DUSKY PIHA,** *Lipaugus fuscocinereus* Cotingidae (114.034) Staffa, 1982, not cataloged, 25 pence

Length: 13 inches, sexes similar, resident. Gray, darker on the wings and tail,


and paler below.


Habitat: montane forest.

Range: Colombia to northern Peru.

Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of the

Birds of the World, Volume 9.





## BLACK-CAPPED TODY-FLYCATCHER, Todirostrum nigriceps

Tyrannidae (116.171)

Bernera, 1982, not cataloged, 1 pound

Length: 3 to 4 inches, sexes alike, resident. Yellowish-olive above and yellow below with a black head and white throat.

Habitat: forest canopy and adjacent clearings.

Range: Eastern Costa Rica to eastern Venezuela and eastern Ecuador. Reference: del Hoyo, J., A. Elliott and J. Sargatal, Eds. Handbook of

the Birds of the World, Volume 9.

[to be continued]